

STUDY OF LOW VOLTAGE ELECTROPORATION TECHNIQUE ON HeLa
CELL TREATED WITH NEOLAMARCKIA CADAMBA EXTRACTS

DAING NOOR FARHANAH BINTI MOHAMAD ISHAK

A thesis submitted in
fulfillment of the requirements for the award of
Degree of Master of Electrical Engineering

Faculty of Electrical and Electronic Engineering
Universiti Tun Hussein Onn Malaysia

MAY 2019

ACKNOWLEDGEMENT

I would like to take this opportunity to express my greatest gratitude to Almighty Allah, for His help and guidance during the course of life and the moment of truth.

I also would like to express my appreciation and sincere gratitude to my supervisor, Associate Professor Ts. Dr. Muhammad Mahadi bin Abdul Jamil and my co supervisor Dr. Mohamad Nazib bin Adon for their continuous support, encouragement and endless patience towards completing this Master research. I am also thankful for their aspiring guidance, invaluable constructive criticism and friendly advice throughout the project work. I am sincerely grateful to both of them for sharing their truthful and illuminating views on a number of issues related to this project.

Further I also would like to convey appreciation to Dr. Salehudin bin Hamdan and Ms. Sayang binti Baba from the Faculty of Bioscience and Medical Engineering, Universiti Teknologi Malaysia and to Dr Ridhwan bin Abd Wahab @ Abdul Wahab from the Department of Biomedical Science, Kulliyyah of Allied Health Sciences, International Islamic University Malaysia, who provided me with the necessary material and equipment for my Master research project and also for their support and guidance.

I would also like to express my appreciation to my beloved husband, Mohd Zulfadli bin Ramli and for his tremendous patience, effort, support and encouragement during my Master research which is highly appreciated.

I would also like to thank both my beloved parents, Mohamad Ishak bin Daing Manaik and Normah binti Noordin and also my in-laws Ramli bin Deraman and Siti Norhayati binti Mohd Daud for their support and encouragement during my master research.

I also would like to express my appreciation to my friends, Azra Hazwanie binti Azizulkarim for their support, encouragement and advice during my masters. Finally, my appreciation to UTHM, for their sponsorship of my studies via Contract

Grant (Vote: U775) for 10 months and also to all for having directly or indirectly contributed towards for the accomplishment of this Master research.



ABSTRACT

Electroporation is a process of creating pores on cell by exposing the cell with electric field. This process increases the permeability of the cell. Neolamarckia Cadamba is a natural source that is widely used in the medical arena due to its pharmacological properties. Neolamarckia Cadamba known as *kelempayang* to Malaysian people. Its mainly planted in the states of Pahang, Perak, Sabah and Sarawak. Electrochemotherapy is a combination of the electroporation process and chemotherapeutic drugs. The Standard electric field parameters for Electrochemotherapy (SEFPECT) are 1000V/cm of pulse amplitude, 100 μ s of pulse duration, 8 numbers of pulses and 1Hz of frequency. Electrochemotherapy process causes transient burns to patient due to the intensity of pulse amplitude used, which is 1000V/cm. The use of chemotherapeutic various drugs gives side effect to the patient. To overcome the problem, this research suggests to lower the pulse amplitude and longer pulse duration plus combining natural extract of Neolamarckia Cadamba Leaves. In this study, the combination of electroporation process and natural extract is conducted in vitro for 48 hours. The cell used is cervical cancer cell (HeLa cell) and the natural extract from Neolamarckia Cadamba Leaves. The experiment was started to determine the best parameter of pulse amplitude and pulse duration that gives lower viability percentage of HeLa cell. The pulse amplitude started with a 200V/cm, 400V/cm, 600V/cm, 800V/cm and 1000V/cm and the pulse duration at 100 μ s, 500 μ s, 1ms, 2ms and 5ms and the result revealed that the best parameter at 600V/cm of pulse amplitude and 5ms of pulse duration. Then the IC₅₀ value of Neolamarckia Cadamba Leaves extract were tested and found to be at 40 μ g/ml. The findings of these studies have shown positive respond on the combination of these two methods. This was proven by the reduction in the percentage of anti proliferation of HeLa cell where with electroporation 30.08% cell can survive and by combining electroporation and extact of Neolamarckia Cadamba leaves only 2.12% HeLa cell can survive.

ABSTRAK

Elektroporasi adalah proses mewujudkan liang-liang pada sel dengan mendedahkan sel dengan medan elektrik. Proses ini meningkatkan kebolehtelapan sel. *Neolamarckia Cadamba* adalah sumber semulajadi yang banyak digunakan di arena perubatan kerana sifat farmakologinya. *Neolamarckia Cadamba* dikenali sebagai kelempayang oleh rakyat Malaysia. Ia kebanyakannya ditanam di negeri Pahang, Perak, Sabah dan Sarawak. Elektrokimoterapi adalah gabungan dari proses elektroporasi dan ubat kemoterapi. Parameter bidang medan elektrik untuk Elektrokimoterapi (SEFPECT) adalah 1000V/cm nadi amplitud, 100 μ s nadi denyut, 8 nadi dan 1Hz frekuensi. Proses elektrokimoterapi menyebabkan kesan terbakar sementara kepada pesakit kerana intensiti nadi amplitud yang digunakan, iaitu 1000V/cm. Penggunaan pelbagai ubat kimoterapi memberi kesan sampingan kepada pesakit. Untuk mengatasi masalah ini adalah dengan cara menurunkan nadi amplitud dan panjangkan tempoh nadi denyut serta menggunakan ekstrak sumber semulajadi daun *Neolamarckia Cadamba*. Dalam kajian ini, gabungan proses elektroporasi dan ekstrak semula jadi dilakukan secara in vitro selama 48 jam. Sel yang digunakan adalah sel kanser serviks (sel HeLa) dan ekstrak semula jadi dari daun *Neolamarckia Cadamba*. Eksperimen ini bermula dengan menentukan parameter nadi amplitud dan nadi denyut yang terbaik memberikan peratusan daya tahan yang lebih rendah kepada sel HeLa. Nadi amplitud bermula dengan 200V/cm, 400V/cm, 600V/cm, 800V/cm dan 1000V/cm dan nadi denyut pada 100 μ s, 500 μ s, 1ms, 2ms dan 5ms dan hasilnya menunjukkan bahawa parameter terbaik pada 600V/cm nadi amplitud dan 5ms nadi denyut. Kemudian nilai IC₅₀ yang ditunjukkan oleh ekstrak daun *Neolamarckia Cadamba* didapati berada pada 40 μ g/ml. Penemuan kajian ini telah menunjukkan tindak balas positif mengenai gabungan kedua-dua kaedah, ini dibuktikan dengan penurunan kadar peratusan percambahan sel HeLa apabila didedahkan medan elektrik hanya 30.08% sel HeLa Bertahan hidup manakala apabila sel HeLa didedahkan dengan gabungan medan elektrik dan ekstrak hanya 2.12% sel hidup.

TABLE OF CONTENTS

TITLE	i
ACKNOWLEDGEMENT	vi
ABSTRACT	viii
ABSTRAK	ix
TABLE OF CONTENTS	x
LIST OF TABLES	xv
LIST OF FIGURES	xvi
LIST OF SYMBOLS AND ABBREVIATIONS	xviii
 CHAPTER 1 INTRODUCTION	 1
1.1. Overview	1
1.2. Background of Research	1
1.3. Problem Statement	3
1.4. Aim	4
1.5. Objectives	4
1.6. Scopes of Study	5
 CHAPTER 2 LITERATURE REVIEW	 6
2.1. Overview	6
2.2. Living Cell	6
2.3. HeLa Cells (Human Cervical Cancer Cells)	8
2.4. Electroporation	10
2.4.1 Membrane Electroporation	10
2.4.2 Application of Electroporation	11
2.4.1 Electric field parameter used by other researcher	12

2.5.	Electrochemotherapy	13
2.6.	Anti-Cancer Agents	14
2.7.	Neolamarckia Cadamba	15
2.7.1	History and Uses of Neolamarckia Cadamba	17
2.7.2	The use of Neolamarckia Cadamba Leaves	19
2.7.3	Phytochemistry found in Neolamarckia Cadamba Leaves and its pharmacological properties	20
2.7.4	Pharmacological effect of Neolamarckia Cadamba: Anti-Cancer or Anti-Tumor	21
2.7.	Summary	21

CHAPTER 3 MATERIALS AND METHODS 24

3.1.	Overview	24
3.2.	Material and Workstation	25
3.2.1	Reagents	25
3.2.2	Equipments	28
3.3.	HeLa Cell Culture Procedure	34
3.3.1	HeLa cell culture result	38
3.4.	HeLa Cell Changing Media Procedure	40
3.5.	HeLa Cell Counting Procedure	40
3.6.	Electroporation Procedure	42
3.7.	Neolamarckia Cadamba Leaves Extract Procedures	45
3.8.	Inhibitory Concentration (IC ₅₀) Determination	47

CHAPTER 4	RESPONSE OF HELA CELL AFTER ELECTROPORATION PROCESS	48
4.1.	Overview	48
4.2.	Electroporation	48
4.3.	Material and Method	49
4.3.1	Equipment	49
4.3.2	Electroporation Procedure	49
4.4.	Effect of electroporation process on HeLa cell	49
4.5.	Comparison between length of HeLa cell without electric field exposed and length of HeLa cell with electric field exposed	52
CHAPTER 5	LOW VOLTAGE PULSE ELECTRIC FIELD EXPOSURE ON HELA CELL	54
5.1.	Overview	54
5.2.	Introduction	54
5.3.	Material and Method	55
5.3.1.	HeLa Cell Culture	55
5.3.2.	Electroporation Method	55
5.3.3.	HeLa Cell Counting Procedure	56
5.4.	Electroporation result for different pulse duration effect on HeLa cell	56
5.5.	Electroporation result for different range of pulse amplitude effect on HeLa	58
5.6.	Electroporation result effect of HeLa cell induced with electric field of 600v/cm and 5ms compared with HeLa cell not induced with electric field	59
5.7.	Summary	62
CHAPTER 6	NATURAL EXTRACT OF NEOLAMARCKIA CADAMBA LEAVES EFFECT ON HELA CELL	64
6.1.	Overview	64
6.2.	Introduction	64

6.3.	Material and Method	66
6.3.1.	HeLa Cell Culture	66
6.3.2.	Neolamarckia Cadamba Leaves Extract Procedure	66
6.3.3.	HeLa Cell Counting Procedure	66
6.3.4.	Inhibitory Concentration (IC ₅₀) Determination	66
6.4.	Different Concentration of Neolamarckia Cadamba Leaves natural extract effect on HeLa cell	67
6.5.	Comparison Result Between HeLa Cell Without Natural Extract Of Neolamarckia Cadamba Leaves And HeLa Cell with Natural Extract Of Neolamarckia Cadamba Leaves	69
6.6.	Summary	72

CHAPTER 7 COMBINED EFFECT OF PULSE ELECTRIC FIELD AND NATURAL EXTRACT OF NEOLAMARCKIA CADAMBA LEAVES ON HELA CELL VIABILITY

7.1.	Overview	74
7.2.	Introduction	75
7.3.	Material and Method	76
7.3.1.	HeLa Cell Culture	76
7.3.2.	Electroporation Method	76
7.3.3.	Neolamarckia Cadamba Leaves Extract Preparation Ptocedure	76
7.3.4.	HeLa Cell Counting Procedure	76
7.4.	Percentage result for HeLa cell proliferation rate, HeLa cell with Electroporation, HeLa cell with Neolamarckia Cadamba Leaves and HeLa cell combined with electroporation process and Neolamarckia Cadamba Leaves extract.	77

7.5.	Untreated HeLa cell and Treated HeLa cell with 40 μ gram of natural extract Neolamarckia Cadamba leaves and electroporation process (600v/cm of voltage and 5ms pulse duration)	79
7.6.	Summary	82

CHAPTER 8	CONCLUSIONS & RECOMMENDATIONS	84
------------------	--	-----------

REFERENCES	88
-------------------	-----------

LIST OF PUBLICATIONS AND COPYRIGHTS	100
--	------------

VITA	102
-------------	------------



PTTA UTHM
PERPUSTAKAAN TUNKU TUN AMINAH

LIST OF TABLES

2.1	Table shows summary of parameter used from previous research done by other researcher	22
2.2	Table shows summary of Neolamarckia Cadamba used from previous research done by other researcher	23
5.1	Percentage means, standard deviation and standard error of electroporation result for different pulse duration effect on HeLa cell	57
5.2	Percentage means, standard deviation and standard error for electroporation effect on HeLa cell with different range of pulse amplitude	59
5.3	Percentage means, standard deviation and standard error for electroporation result effect of HeLa cell induced with electric field of 600v/cm and 5ms compared with HeLa cell not induced with electric field	61
6.1	Percentage means, Standard Deviation and Standard Error for different Concentration of Neolamarckia Cadamba leaves natural extract effect on HeLa cell	68
6.2	Percentage means, Standard Deviation and Standard Error for comparison result between HeLa cell without natural extract of Neolamarckia Cadamba Leaves and HeLa cell with natural extract of NeolamarckiaCadamba Leaves	71
7.1	Percentage means, standard deviation and standard error for HeLa cell proliferation rate, HeLa cell with Electroporation, HeLa cell with Neolamarckia Cadamba Leaves and HeLa cell combined with electroporation process and Neolamarckia Cadamba Leaves extract	78
7.2	Percentage means, standard deviation and standard error for Untreated HeLa cell and treated HeLa cell with 40µgram of natural extract Neolamarckia Cadamba leaves and electroporation process (600v/cm voltage and 5ms pulse duration)	81

LIST OF FIGURES

2.1	Prokaryotic cell	7
2.2	Eukaryotic cell	8
2.3	Human Cervical Cancer Cell (HeLa Cell) scale bar 50µm	9
2.4	Neolamarckia Cadamba Tree	16
2.5	Neolamarckia Cadamba Fruit	17
2.6	Neolamarckia Cadamba Leaves	18
2.7	Neolamarckia Cadamba Bark	19
3.1	Flowchart of overall experiment	24
3.2	Rosewell Park Memorial Institute (RPMI) medium 1640 (Gibco) or RPMI Medium	26
3.3	Fetal Bovine Serum (FBS) 10270 (Gibco)	26
3.4	Penicilin Streptomycin Antibiotics (Gibco)	27
3.5	Phosphate Buffered Saline (PBS) PH7.4 (Gibco)	27
3.6	trypLE Express trypsin (Gibco)	28
3.7	Conical Flask	28
3.8	Culture Flask	29
3.9	Pipette	29
3.10	Biological Safety Cabinet	30
3.11	CO ₂ Incubator	30
3.12	Microscope Nikon ECLIPSE TS100	31
3.13	Waterbath	32
3.14	Centrifuge MPW-56	32
3.15	6 well-plated dishes	33
3.16	ECM 830 Electroporation	33
3.17	Cuvette 4mm gap	34
3.18	Hemocytometer	34
3.19	HeLa cell after 6-7 days subculture	35
3.20	HeLa cell after it is completely detached from the surface of culture flask	36
3.21	Culture Procedure	37

3.22	HeLa cell culture picture taken every 6 hours	39
3.23	Hemocytometer glass	41
3.24	grid lines of the hemocytometer	42
3.25	Electroporation procedure	44
3.26	Neolamarckia Cadamba Leaves Extract Procedure	46
4.1	Picture of HeLa cell length measured using DinoCapture 2.0 software	50
4.2	Picture showing HeLa cell length result after electric field exposed to HeLa cell	51
4.3	A) HeLa cell without electric field exposed B) HeLa cell with electric field exposed	53
5.1	Graph showing the electroporation result for different pulse duration effect on HeLa cell	57
5.2	Graph showing the electroporation result for different range of pulse amplitude effect on HeLa cell	59
5.3	Graph showing the electroporation result effect of HeLa cell induced with electric field of 600v/cm and 5ms compared with HeLa cell not induced with electric field	62
6.1	Neolamarckia Cadamba Leaves	65
6.2	Graph showing result for different Concentration of Neolamarckia Cadamba Leaves natural extract effect on HeLa cell	69
6.3	Graph showing comparison result between HeLa cell without natural extract of Neolamarckia Cadamba Leaves and HeLa cell with natural extract of Neolamarckia Cadamba Leaves	72
7.1	Graph for HeLa cell proliferation rate, HeLa cell with Electroporation, HeLa cell with Neolamarckia Cadamba Leaves and HeLa cell combined with electroporation process and Neolamarckia Cadamba Leaves extract	79
7.2	Graph shows the result for untreated and treated HeLa cell with 40 μ gram of natural extract Neolamarckia Cadamba leaves and electroporation process (600v/cm voltage and 5ms pulse duration)	82

LIST OF SYMBOL AND ABBREVIATIONS

- ECM 830* - Square Wave Pulse Generator
- EP* - Electroporation
- HeLa* - Cervical Cancer Cell
- SEFPECT* - Standard electric field parameters for Electrochemotherapy



CHAPTER 1

INTRODUCTION

1.1. Overview

This chapter consists of background of research followed by problem statement, aim, objectives and scope of study.

1.2. Background of Research

HeLa cells are human cervical cancer cells (the cervix is found at the top of the vagina and the entrance of the uterus). The cells were first cultured or grown in the laboratory in 1951 and were the first type of human cancer cell to be cultured continuously for experiments. There are many different strains (subtypes) of HeLa cells that are now used (J. Zhang et al., 2015).

Electroporation or electropermeabilization is a molecular biology technique in which an electrical field is applied to cells in order to increase the permeability of the cell membrane, allowing chemicals, drugs, or DNA to be introduced into the cell (C. Rosazza et al., 2016). In molecular biology, the process of electroporation is often used to transform bacteria, yeast, or plant protoplasts by introducing new coding DNA. When bacteria and plasmids are mixed together, the plasmids can be transferred into the bacteria after electroporation. Several hundred volts across a

distance of several millimeters are typically used in this process. Afterwhich, the cells have to be handled carefully until they are able to divide and produce cells that contains reproduced plasmids. This process is approximately ten times more effective than chemical transformation (A. Natarajan et al., 2012).

Electroporation is also highly efficient for the introduction of foreign genes into tissue culture cells, especially mammalian cells. For example, it is used in the process of producing knockout mice, as well as in tumor treatment, gene therapy, and cell-based therapy (C. R. Tracy et al., 2010). The process of introducing foreign DNA into eukaryotic cells is known as transfection. Electroporation is highly effective for transfecting cells in suspension using electroporation cuvettes. Electroporation has proven efficient for use on tissues in vivo, for in utero applications as well as in ovo transfection. Adherent cells can also be transfected using electroporation, providing researchers with an alternative of trypsinizing their cells prior to transfection (A. Rizk & B. M. Rabie, 2013).

Compound Extract Test as Anti-Proliferation (Neolamarckia Cadamba Leaves) also well-known as “Anthocephalus Cadamba”, is a precious plant in Ayurvedic medicine. This Rubiaceae family tree, is commonly called as Kadam by the Indians and Kalempayan by Malaysians which it grows naturally in Australia, India, China, Malaysia, Indonesia, Philipines, Vietnam, and Singapore. Moreover, Neolamarckia Cadamba tree can grow up to 45 m tall, with a diameter up to 100-160 cm but normally less. Neolamarckia Cadamba has an umbrella shape crown and tiers arrangement of branches (C. P. Khare, 2007)

Importantly, Neolamarckia Cadamba is selected as one of the timber species for forest rehabilitation projects in Malaysia, due to its short rotation period and early commercial returns within 8-10 years. Whilst in India, the leaves are traditionally used in treating illnesses. It has been mentioned and described in many Indian medicine for the treatment of anemia, fever, uterine and liver complications, diabetes, diarrhea and much more. It also has antioxidant, anti-inflammatory also wound healing, anti-tumor, and anti-cancer properties. (A. Kumar et al., 2015).

1.3. Problem Statement

Statistically, cancer is among the leading causes of morbidity and mortality worldwide with approximate 18 million cancer cases recorded based on World Health Organization in 2018. This number is expected to increase by about 63% over the next 2 decades. In Malaysia, the numbers of cancer cases are approximately at 43 thousand recorded in 2018. This number is expected to increase by 47% over the next 2 decades. In Malaysia the most common cancer diagnosed in 2018 are breast, cervical, prostate, lung and colorectal cancer (F. Bray, et al., 2018).

Cancer can start at any part of the body. It starts when cell growth is out of control and crowds out the normal cell, thus making it hard for the body to function as it should. Cancer cells spread to other parts of the body. For instance, cancer cells in the lung can spread to the bones and grow. When cancer cells spreads, it is called metastasis. Some cancers cells grow and spread fast and others grow slowly. Most cancer form a lump called a tumor or a growth. But not all lumps are cancerous. Lumps that are cancerous are called malignant which have the potential to be dangerous (G. Cooper & R. E. Hausman, 2000). An alternative cancer chemoprevention is a medical intervention that aims to reduce the incidence and burden of cancer via the development of agents to prevent, reverse and delay the process of carcinogenesis (W. P. Steward & K. Brown, 2013).

The current chemotherapy side effect on human depends on the cancer type, drug taken and dosage. The treatment is not controllable (J. B. Wolinsky, et al., 2012). This study suggests an alternative treatment by used natural extract of *Neolamarckia Cadamba* leaves. This natural extract has phenolic found inside the leaves that have anti-cancer properties.

1.4. Aim

The aim of this research is to investigate the effect of electroporation and natural extracts of Neolamarckia Cadamba Leaves on HeLa cell cultured

1.5. Objectives

This study embarks on the following objectives:

- a) To identify electric field parameter due to Electroporation process towards HeLa cell.
- b) To analyze the effect of Neolamarckia Cadamba Leaves on HeLa cell
- c) To evaluate the combined effect of Electric Field and Neolamarckia Cadamba Leaves on HeLa cell as anti-cancer mechanism.

1.6. Scope of study

In order to achieve the objectives of this research, the following scope of work were identified as followed:

a) HeLa cell lines were acquired and cultured in a physiological environment similar to mammalian cell. The cells were maintained at temperature of 37 °C similar to human temperature and humidified environment at 5% CO₂ in order to maintain the pH of the medium as in the human blood plasma (J. T. Lee, et al., 2011).

b) Investigation of the best electric field parameter effect on HeLa cell lines proliferation in vitro followed by the standard electric field parameter for electrochemotherapy (SEFPECT) not more than 1000V/cm of pulse amplitude using the following parameter:

Pulse Amplitude: 200V/cm, 400V/cm, 600V/cm, 800V/cm and 1000V/cm

Pulse Duration: 100µs, 500µs, 1ms, 2ms and 5ms

c) Investigation of anti-cancer properties and determination 50% Inhibitory Concentration (IC₅₀) of Neolamarckia Cadamba leaves effect on HeLa cell lines proliferation, this is by considered the following range of concentration: 1µg/ml, 5µg/ml, 10µg/ml, 20µg/ml, 30µg/ml, 40µg/ml, 50µg/ml, 60µg/ml, 70µg/ml, 80µg/ml, 90µg/ml and 100µg/ml

d) Analyse the Combined effect of electric field and Neolamarckia Cadamba leaves extract on HeLa cell lines

e) Experiment to be repeated minimum thrice or more to ensure the reading is accurate and to get an average value.

CHAPTER 2

LITERATURE REVIEW

2.1. Overview

This chapter focuses on previous research done by other researchers that is similar and related to this research. This chapter is important to give an idea, guide and knowledge to the researcher on the commencement of the experiment.

2.2. Living Cell

Based on Mariana (2007) cells are called “Building blocks of life”. The author described that cells are the functional and structural unit of all organisms that are living in this world. Cells are divided into two types that are prokaryotic cells and eukaryotic cells. The characteristics of prokaryotic cells can be described by the size that is usually less than 5µm and smaller compared to eukaryotic cells. This type of cell does not have a nucleus. Prokaryotic cells mass typically is around 1 nano gram. Examples of prokaryotic cells are bacteria and archaea. Figure 2.1 showed the structure of the prokaryotic cell.




Figure. 2.1 Prokaryotic cell

(<http://legacy.hopkinsville.kctcs.edu/instructors/Jason-Arnold/VLI/Module%202/m2cellstructure/m2cellstructure4.html> and <http://7marshscience.blogspot.com/2015/09/>)

For a eukaryotic cell, the characteristics are opposed to a prokaryotic cell. Eukaryotic cells usually have more complex structures. The structures of eukaryotic cells are called organelles. Organelles contain many structures such as the nucleus, endoplasmic reticulum that consists of rough endoplasmic reticulum (contain ribosomes) and smooth endoplasmic reticulum (does not contain ribosomes), Golgi apparatus, lysosomes, mitochondria, chloroplast (found inside plant cells), vesicles, vacuoles and many more. Animal cell, plant cell, fungi, and protista are examples of eukaryotic cells. Figure 2.2 shows the structure of the eukaryotic cell.




Figure. 2.2 Eukaryotic cell

(<http://legacy.hopkinsville.kctcs.edu/instructors/Jason-Arnold/VLI/Module%202/m2cellstructure/m2cellstructure4.html> and <http://7marshscience.blogspot.com/2015/09/>)

2.3. HeLa Cells (Human Cervical Cancer Cells)

HeLa cells are one of the examples of eukaryotic cell type. It is derived from human cervical cancer cells. HeLa is named based on cancer patient name Henrietta Lacks. This cell was taken from Henrietta Lacks who passed away on 4 October 1951 after fighting for eight months from her cervical cancer disease. HeLa cells were first cultured in 1951 in a laboratory (J. Zhang et al., 2015). It has been continuously grown as first human cells known. HeLa cells have a high capability of division and growing indefinitely and continuously as long as the environment of the cell is still

suitable for HeLa cells to grow and adapt. This characteristic made HeLa cells easily immortalize literally.

For a decade in cancer research area, many scientists and researchers explored and studied the HeLa cells characteristics due to its availability and easiness to find because it is cultured continuously in the laboratory. HeLa cells are very tough and adaptable cell. It can survive in any ambient where other cells cannot and die easily. HeLa cell is also a very potent cell that is able to contaminate another cell. Figure 2.3 showed the structure of HeLa cell taken during experiment.



Figure. 2.3 Human Cervical Cancer Cell (HeLa Cell) scale bar 50μm

2.4. Electroporation

Electroporation is the method of external application of electric field to the cell membrane to increase the permeability of the cell membrane. This method is widely used in medicines, food processing and pharmaceuticals (M. Yarmush et al., 2014). Electroporation also is a method to create pores on cell membrane to allow any substances to enter the cell easily (N. J. Yang & M. J. Hinner, 2014).

2.4.1. Membrane Electroporation

R. Stampfli (1958), stated that our bodies are built from trillions of biological cells enveloped by the plasma membrane. The plasma membrane is largely composed of a double layer (bilayer) of lipids. Its size is about 5nm. The inside of the membrane is partly gel and partly liquid. It functions as a protector and separator of the cell from its surroundings environment. It is very stable and reliable. The lipid bilayer contains different types of proteins which act as pumps and channel for transporting purpose. It can transport specific molecules that want to across the plasma membrane. Plasma membrane can be electrically be viewed as a sheet of thin insulator surrounded by electrolyte aqueous solutions on both sides.

Based on A. J. H. Sale & W. A. Hamilton (1967), when strong electroporation is exposed to the membrane, it will undergo electrical breakdown that allows it to be permeable to other molecules to freely cross it. This process is called the electroporation of the membrane. If the exposure of electric field is sufficiently short, then the membrane can effectively undergo rapid recovery for the cell to remain viable. It is called reversible electroporation. Otherwise, it is called irreversible electroporation (E. Neumann & K. Rosenheck, 1972).

Electroporation becomes a useful tool in biotechnology and medicine area. As for irreversible electroporation, it becomes popular for electrochemotherapy where the method is by introducing drugs into tumor cells (G. Sersa et al., 2008). Another

REFERENCES

- A. Agarwal, I. Zudans, E. A. Weber, J. Olofsson, O. Orwar, S. G. Weber, "Effect of cell size and shape on single-cell electroporation", *Analytical chemistry* 79(10):3589-3596 (2007).
- A. Dwevedi, K. Sharma, & Y. K. Sharma, "Cadamba: A miraculous tree having enormous pharmacological implications," *Pharmacognosy Reviews*, 9(18), 107–113, (2015).
- A. Gupta, M. Anand, S. Yadav, J. Gautam, "Phytochemical studies and antioxidant activity of different leaves extracts of A. Cadamba," *International Journal of Futuristic Science Engineering and Technology*, Vol 1, ISSN 2320 – 4486, (2013).
- A. J. H. Sale, & W. A. Hamilton, "Effects of high electric fields on microorganisms killing of bacteria and yeasts," *Biochem. Biophys. Acta* 148, 781-788, (1967).
- A. Kumar, S. R. Chowdhury, K. K. Jatte, T. Chakrabarti, H. K. Majumder, T. Jha, S. Mukhopadhyay, "Anthocephaline, a new indole alkaloid and cadambine, a potent inhibitor of DNA topoisomerase IB of *Leishmania donovani* (LdTOP1LS), isolated from *Anthocephalus Cadamba*," *Nat. Prod. Commun.* 10, 297–299, (2015).
- A. M. Gonzalez-Augulo, F. Morales-Vasquez, and G. N. Hortobagyi, "Overview of resistance to systemic therapy in patients with breast cancer", Ch 1, pp. 1-22, in *Breast Cancer Chemosensitivity*, D. Yu and M-C Hung (Eds), Springer Science Business Media, LLC, Landes Bioscience, New York, USA, (2007).
- A. M. Lebar, G. Sersa, R. Kranjc, A. Groselj, & D. Miklavcic, "Optimisation of pulse parameters in vitro for in vivo electrochemotherapy. Anticancer Research," 22(3), pp. 731-1736. (2002).

- A. Natarajan, L. Campana, and R. Sundararajan, "Electrically-Enhanced Delivery of Drugs and Conjugates for Cancer Treatment," *Antibody-Mediated Drug Delivery Systems*, pp. 129–143, (2012).
- A. Rizk and B. M. Rabie, "Electroporation for Transfection and Differentiation of Dental Pulp Stem Cells," *BioResearch Open Access*, vol. 2, no. 2, pp. 155–162, (2013).
- A. V. Titomirov, S. Sukharev, et al., "In vivo electroporation and stable transformation of skin cells of newborn mice by plasmid DNA," *Biochim. Biophys. Acta* 1088, 131-134. (1991).
- B. D. Sukhendu, P. Dietmar, Rabussay, G. Widera, and G. A. Hofmann, "Medical Applications of Electroporation", *Ieee Transactions On Plasma Science*, Vol. 28, No. 1, February (2000).
- C. De. Martel, J. Ferlay, et al., "Global burden of cancers attributable to infections in 2008: a review and synthetic analysis," *The Lancet Oncology* 2012;13: 607-615, (2012)
- C. P. Khare, "Indian Herbal Remedies: Rational Western Therapy, Ayurvedic and Other Traditional Usage, Botany," *Springer Science & Business Media*, NewYork, USA, pp.66–67, (2011).
- C. P. Khare, "Indian Medicinal Plants: An Illustrated Dictionary," *Springer*, New York, USA. (2007).
- C. Rosazza, S. Haberl Meglic, A. Zumbusch, M. Rols and D. Miklavcic, "Gene Electrotransfer: A Mechanistic Perspective", *Current Gene Therapy*, vol. 16, no. 2, pp. 98-129, (2016).
- C.R. Tracy, W. Kabbani, J.A. Cadeddu, "Irreversible electroporation: a novel method for renal tissue ablation", *BJU International*, (2010).
- D. A. Dean, T. Ramanathan, D. Machado, and R. Sundararajan, "Electrical impedance spectroscopy study of biological tissues," *Journal of Electrostatics*, vol. 66, no. 3-4, pp. 165-177, (2008).

- D. Ganjewala, N. Tomar, A. K. Gupta “ Phytochemical Composition And Antioxidants Properties of Methanol Extracts of Leaves and Fruits of *Neolamarckia cadamba* (Roxb.)”, *Journal of Biologically Active Products from Nature* , 3(4), 232-240. (2013)
- D. Joker, “*Neolamarckia cadamba* (Roxb.) Bosser (*Anthocephalus chinensis* (Lam.) A. Rich. ex Walp),” (2000)
- D. P. Mishra, M. A. Khan, D. K. Yadav, A. K. Rawat, R. K. Singh, T. Ahamad, M. K. Hussain, M. Saquib, and M. F. Khan, “Monoterpene Indole Alkaloids from *Anthocephalus cadamba* Fruits Exhibiting Anticancer Activity in Human Lung Cancer Cell Line H1299,” *ChemistrySelect*, vol. 3, no. 29, pp. 8468–8472, Jan. (2018).
- D. Patel, V. C. Darji, A. H. Bariya, K. R. Patel, R. N. Sonpal, “Evaluation of anti-fungal activity of *Neolamarckia cadamba* (Roxb.) Bosser leaf and bark extract,” *Int. Res. J. Pharm.* 2, 192–193, (2011)
- E. Neumann and K. Rosenheck, “Permeability changes induced by electric impulses in vesicular membranes,” *J. Membrane Biol.*, vol. 10, pp. 279–290, (1972).
- E. Neumann, M. Schafer-Ridder, et al, “Gene transfer into mouse lyoma Cells by electroporation in high electric fields,” *EMBO J.* 1, 841-845 (1982).
- E. W. Lee, S. Thai, and S. T. Kee, “Irreversible electroporation: A novel image-guided cancer therapy,” *Gut Liver*, vol. 4, pp. S99–S104, (2010).
- F. André and L. M. Mir, “DNA electrotransfer: Its principles and an updated review of its therapeutic applications,” *Gene Ther.*, vol. 11, pp. S33–S42, (2004).
- F. Bray, J. Ferlay, I. Soerjomataram, R. L. Siegel, L. A. Torre, and A. Jemal, “Global cancer statistics 2018: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries,” *CA: A Cancer Journal for Clinicians*, vol. 68, no. 6, pp. 394–424, (2018).
- F. Xiao, J. F. Leary, L. Reece, I. G. Camarillo, M. Sundararajan, R. Sundararajan, “Nanosecond, Low Voltage Electro-Endocrine-Therapy for Breast Cancer

Treatment, 2010 Annual Report Conference on Electrical Insulation and Dielectric Phenomena, (2010)

G. Cooper and R. E. Hausman, *The cell ; a molecular approach*. Oxford: Sinauer Associates, (2000).

G. M. Cragg and D. J. Newman, "Plants as a source of anti-cancer agents," *Journal of Ethnopharmacology*, vol. 100, no. 1-2, pp. 72–79, (2005).

G. Pucihar, L. M. Mir, & D. Miklavcic, The effect of pulse repetition frequency on the uptake into electroporabilized cells in vitro with possible applications in electrochemotherapy. *Bioelectrochemistry*, 57(2), pp. 167-172.(2002).

G. Sersa, "Electrochemotherapy: Animal model work review. In:Electrochemotherapy, electrogenetherapy and transdermal drug delivery," pp. 119-136. Eds., Jaroszeski, M. J., Heller, R., Gilbert, R., Humana Press, Totowa, NJ. (2000).

G. Sersa, D. Miklavcic, "Electrochemotherapy of Tumours", *Journal of Visualized Experiments*, (2008).

G. Sersa, D. Miklavcic, M. Cemazar, Z. Rudolf, G. Pucihar, and M. Snoj, "Electrochemotherapy in treatment of tumours," *Eur. J. Surg. Oncol.*, vol. 34, pp. 232–240, (2008).

G. Sersa, M. Cemazar, & M. Snoj, Electrochemotherapy of tumours. *Curr Oncol*, 16(2), pp. 34-35. (2009)

H. A. Pohl, & J. S. Crane, "Dielectrophoresis of cells," *Biophys. J.*11, 711-727. (1971).

H. B. Mamman, M. M. A. Jamil, M. N. Adon, "Low amplitude pulse electric field for elimination of unpleasant sensation associated with high amplitude electric field for electrochemotherapy," *Journal of Advances in Technology and Engineering Studies*, 3(2): 27-33, (2017).

- H. B. Mamman, M. M. A. Jamil, M. N. Adon, "Optimization of Electric Field Parameters for HT29 Cell Line towards Wound Healing Application," *Indian Journal of Science and Technology*, Vol 9 (46), (2016).
- H. G. L. A. Coster, "Quantitative analysis of the voltage-current relationships of fixed charge membranes and the associated property of "punch-through"," *Biophys. J.* 5, 669-686. (1965).
- H. Gurjar, S. K. Jain, R. Irchhaiya, R. Nandanwar, V. K. Sahu, H. Saraf, "Hypoglycemic effects of methanolic extract of *Anthocephalus cadamba* bark in alloxan induced diabetic rats (RoxB)," *Miq. Int. J. Pharm. Sci. Res.* 1,79–83, (2010).
- H. H. Aung, C. Z. Wang, M. Ni, A. Fishbein, S. R. Mehendale, J. T. Xie, A. Y. Shoyama, C. S. Yuan, "Crocins from *crocus sativus* possesses significant anti-proliferation effects on human colorectal cancer cells," *Exp Oncol.*, 29(3): 175–180. (2007).
- H. Takayamma, S. Tsutsumi, M. Kitajima, D. Santiarworn, B. Liawruangrath, N. Aimi, "Glucosyl-indole alkaloids from *Nauclea cadamba* in Thailand and transformation of 3 α -dihydrocadambine into the indolopyridine alkaloid, 16-carbomethoxynaupholine," *Chem. Pharm. Bull.* 51,232–233, (2003)
- I. Kitagawa, H. Wei, S. Nagao, T. Mahmud, K. Hori., M. Kobayashi, T. Uji, H. Shibuya, "Indonesian Medicinal Plants XIV. Characterization of 3'-O-Caffeoyls- weroside, a new secoiridoid glucoside, and kelampayosides A and B, two new phenolic apioglucosides, from the bark of *Anthocephalus chinensis* (Rubiaceae)," *Chem. Pharm. Bull.* 44,116–127, (1996)
- J. B. Wolinsky, Y. L. Colson, and M. W. Grinstaff, "Local drug delivery strategies for cancer treatment: Gels, nanoparticles, polymeric films, rods, and wafers," *Journal of Controlled Release*, vol. 159, no. 1, pp. 14–26, (2012).
- J. J. M. Landry, P. T. Pyl, T. Rausch, T. Zichner, M. M. Tekkedil, A. M. Stütz, A. Jauch, R. S. Aiyar, G. Pau, N. Delhomme, J. Gagneur, J. O. Korbel, W. Huber, and L. M. Steinmetz, "The Genomic and Transcriptomic Landscape of a HeLa

- Cell Line,” *Genes Genomes Genetics*, vol. 3, no. 8, pp. 1213–1224, Nov. (2013).
- J. L. Hartwell, "Plants used against cancer. A survey." *Lloydia*, 32, 153, 1969. 15, (1969).
- J. M. Crowley, “Electrical breakdown of bimolecular lipid membranes as an electromechanical instability,” *Biophys. J.* 13, 711-724. (1973).
- J. T. Lee, Y. Leng, K. L. Chow, F. Ren, X. Ge, K. Wang, and X. Lu, “Cell culture medium as an alternative to conventional simulated body fluid,” *Acta Biomaterialia*, vol. 7, no. 6, pp. 2615–2622, (2011).
- J. Zhang, J. Jia, L. Zhao, X. Li, Q. Xie, X. Chen, J. Wang, and F. Lu, Down-regulation of microRNA-9 leads to activation of IL-6/Jak/STAT3 pathway through directly targeting IL-6 in HeLa cell. *Molecular Carcinogenesis*, 55(5), pp.732-742, (2015).
- J.C Weaver and Y.A. Chizmadzhev, “Theory of electroporation : A review,” *Bioelectrochem. Bionerg.*, vol.41, pp.135-160, (1996).
- J.C Weaver, “Electroporation of cells and tissues,” *IEEE Trans. Plasma Sci.*, vol.28, pp. 24-33, (2000).
- L. D. Kapoor, “Handbook of Ayurvedic Medicinal Plants: Herbal,” Reference Library. CRC Press, p.43, (2000).
- L. M. Mir, & S. Orlowski, Mechanisms of electrochemotherapy. *Advanced Drug Delivery Reviews*, 35(1), pp. 107-118. (1999).
- L. M. Mir, J. Gehl, G. Sersa, C. G. Collins, J. R. Garbay, V. Billard, & M. Marty, Standard operating procedures of the electrochemotherapy: instructions for the use of bleomycin or cisplatin administered either systemically or locally and electric pulses delivered by the Cliniporator TM by means of invasive or non-invasive electrodes. *European Journal of Cancer Supplements*, 4(11), pp. 14-25. (2006).

- L. M. Mir, M. Belehraddek, et al., "Electrochemotherapy, a novel antitumor treatment: first clinical trial," *C.R. Acad. Sci. Paris* 313, 613-618, (1991).
- L. Miller, J. Leor, and B. Rubinsky, "Cancer cells ablation with irreversible electroporation", *Technology in Cancer Research and Treatment*, Vol. 4, No. 6, pp. 1-7, Dec (2005).
- M. A. Alam, R. Akter, N. Subhan, M. M. Rahman, M. M. Majumder, Nahar, S. D. Sarker, "Anti diarrhoeal property of the hydro ethanolic extract of the Fruits of *Anthocephalus Cadamba*," *Rev. Bras. Farmacogn.* 18, 155–159, (2008).
- M. A. Hassan, M. A. Ferdous, R. Chowdhury, K. A. Khan, "Evaluation of analgesic, anthelmintic and cytotoxic, potential activity of bark of *Anthocephalus Cadamba*," *Int. J. Innov. Pharm. Sci.d Res.*1, 99–107, (2013)
- M. A. Hossain, M. Z. Sultan, A. M. S. Chowdhury, C. M. Hasan, M. A. Rashid, "Preliminary cytotoxicity and antimicrobial investigation of *Anthocaphalus chinensis*," *J. Sci. Res.* 3,689–692, (2011)
- M. A. M. Zaltum, M. N. Adon, and M. M. A. Jamil, "Electroporation effect on the growth of
- M. A. S. Rafshanjani, S. Parvin, M. A. Kader, "Antimicrobial and preliminary cytotoxic effects of ethanol extract and its fractions of *Anthocephalus cadamba* (Roxb.) Miqstem bark. *Int. J. Pharm. Life Sci.*5,4038–4044, (2014)
- M. Arora, "Cell Culture Media: A Review," *Materials and Methods*, vol. 3, (2013).
- M. B. Sporn & N. Suh, "Chemoprevention: an essential approach to controlling cancer," *Nat Rev Cancer.* 2(7): 537-543,(2002).
- M. Chandel, S. Kaur, S. Kumar, "Studies on the genoprotective/antioxidant potential of methanol extract of *Anthocephalus cadamba* (Roxb.)," *Miq. J. Med. Plants Res.*5, 4764–4770. (2011).
- M. Chandel, U. Sharma, N. Kumar, B. Singh, and S. Kaur, "Antioxidant activity and identification of bioactive compounds from leaves of *Anthocephalus cadamba* by ultra–performance liquid chromatography/electrospray ionization

quadrupole time of flight mass spectrometry,” *Asian Pacific Journal of Tropical Medicine*, 5(12), pp.977-985, (2012).

M. Greenwell and P. K. S. M. Rahman, “Medicinal Plants: Their Use in Anticancer Treatment.” *International journal of pharmaceutical sciences and research* 6.10 (2015)

M. Heish, T. Salameh, I. Camarillo and R. Sundararajan, “Irreversible Electroporation Effects: A Drug-Free Treatment for Cancer,” (2011).

M. L. Lucas, L. Heller, et al., “Electro-gene-therapy for the successful treatment of established subcutaneous B16F10 melanoma,” *Mol. Therapy* 5, 668-675, (2002).

M. Morales-de la Peña, P. Elez-Martínez, and O. Martín-Belloso, “Food preservation by pulsed electric fields: An engineering perspective,” *Food Eng. Rev.*, vol. 3, pp. 94–107, (2011).

M. Okino, M. Marumoto, et al., “Electrical impulse chemotherapy for rat solid tumors,” *Proc. Jpn. Cancer Congress*, 46, 420, (1987).

M. P. Rols,, “Electropermeabilization, a physical method for the delivery of therapeutic molecules into cells”, *Biochimica et Biophysica Acta*, 1758, 423, (2006).

M. Sack, J. Sigler, S. Frenzel, C. Eing, J. Arnold, T. Michelberger, W. Frey, F. Attmann, L. Stukenbrock, and G. Müller, “Research on industrial-scale electroporation devices fostering the extraction of substances from biological tissue,” *Food Eng. Rev.*, vol. 2, pp. 147–156, (2010).

M. Yarmush, A. Golberg, G. Serša, T. Kotnik and D. Miklavčič, "Electroporation-Based Technologies for Medicine: Principles, Applications, and Challenges", *Annual Review of Biomedical Engineering*, vol. 16, no. 1, pp. 295-320, (2014).

- Min Rui, Yingyu Chen, et al, "Transfer of anti-TFAR19 monoclonal antibody into HeLa cell by in situ electroporation can inhibit the apoptosis", 71, 1771–1778, (2002).
- N. A. A. Rahman, M. M. A. Jamil, "Investigation of Pulsed Electric Field on Cancer Cell Cultured on Patterned Surface," 6th IEEE International Conference on Control System, Computing and Engineering, (2016).
- N. Dolai, I. Karmakar, K. R. B. Suresh, B. Kar, A. Bala, P. K. Halder, "Evaluation of antitumor activity and in vivo antioxidant status of Anthocephalus Cadamba on Ehrlich ascites carcinoma treated mice," J.Ethnopharmacol.142,865–870, (2012)
- N. J. Yang and M. J. Hinner, "Getting Across the Cell Membrane: An Overview for Small Molecules, Peptides, and Proteins," Site-Specific Protein Labeling Methods in Molecular Biology, pp. 29–53, Aug. (2014).
- N. P. Julie, T. Hop, T. Hung, A. P. Tuyet, D. Christiane, F. Jeremy, H. T. Tinh, C. Philippe, B. Bernard, G. Philippe, "Antimalarial and cytotoxic activities of ethnopharmacologically selected medicinal plants from South Vietnam," J.Ethnopharmacol. 109,417–427, (2007).
- N. Pavselj, & D. Miklavcic, Numerical modeling in electroporation-based biomedical applications. Radiology and Oncology, 42(3), pp. 159-168, (2008).
- Nordahlia, As & Lim, Seng & Husain, Hamdan & Uyup, Mohd Khairun Anwar. Tb54_Wood Properties Of Selected Plantation Species_Teak Kelempayan Binuang And Batai. (2014).
- P. Nagakannan, B. D. Shivasharan, V. P. Veerapur, B. S. Thippeswamy, "Sedative and antiepileptic effects of Anthocephalus cadamba Roxb.in mice and rats," Indian J.Pharmacol.43,699–702, (2011)
- P. Nagar, K. Diwakar, A. Nagar, S. Gullaiya, V. Dubey, V. Singh, "A pre-liminary study of the effects of petroleum ether and alcoholic bark extract of Anthocephalus cadamba (Roxb.) Miq. In acute and chronic in-vivo inflammatory rat models," Int.Res.J.Pharm.3,217–221, (2012)

- R. Heller, M. Jaroszeski, et al., "In Vivo Gene Electroinjection and Expression in Rat Liver," Fed. Europ. Biochem. Soc. (FEBS) Letters 389, 225-228, (1996).
- R. Heller, R. Gilbert, et al., "Clinical trials for solid tumors using electrochemotherapy. In: Electrochemotherapy, electrogenotherapy and transdermal drug delivery," pp. 137-156. (2000).
- R. P. Mishra, "A Comparative Study and Extract Optimization for Antimicrobial Properties of Different Parts of Anthocephalus Cadamba," ISSN 2046-1690, (2013).
- R. Stampfli, "Reversible electrical breakdown of the excitable membrane of a Ranvier node," An. Acad. Brasil. Ciens., vol. 30, pp. 57–63, (1958).
- R. V. Davalos, L. M. Mir, and B. Rubinsky, "Tissue ablation with irreversible electroporation," Annals of Biomedical Engineering, vol. 33, no. 2, pp. 223–231, Feb. (2005).
- R. Ziv, Y. Steinhardt, G. Pelled, D. Gazit, and B. Rubinsky, "Micro-electroporation of mesenchymal stem cells with alternating electrical current pulses," Biomedical Microdevices, vol. 11, no. 1, pp. 95–101, (2008).
- S. G. Razafimandimbison, ".A systematic crevision of Breonia (Rubiaceae-Nau- cleaeae)", Ann. Missouri Bot.Gard.89,1–37.(2002).
- S. Li, X. Zhang, et al., "Regression of tumor growth and induction of long-term antitumor memory by interleukin 12 electro-gene therapy," J. Natl. Cancer Inst. 94, 762-768. (2002).
- S. Mondal, G. K. Dash, S. Acharyya, "Analgesic, anti-inflammatory and an-tipyretic studies of Neolamarckia cadamba barks," J.Pharm.Res.2, 1133–1136, (2009).
- S. P. Umachigi, G. S. Kumar, K. N. Jayaveera, D. V. K. Kishore, K. C. K. Ashok, R, Dhanpal, "Anti microbial, wound healing and antioxidant activities of Anthoce-phalus cadamba", (2007)

- S. Rashid, A. Unyayar, M. Mazmanci, S. Mckeown, I. Banat, and J. Worthington, "A study of anti-cancer effects of *Funalia trogii* in vitro and in vivo," *Food and Chemical Toxicology*, vol. 49, no. 7, pp. 1477–1483, (2011) .
- S. S. Handa, R. P. Borris, G. A. Cordell, J. D. Phillipson, "NMRspectral analysis of cadambine from *Anthocephalus chinensis*," *J. Nat Prod.* 46,325–330, (1983).
- S. Singh, M. P. S. Ishar, A. K. Saxena, A. Kaur, "Cytotoxic effect of *Anthocephalus cadamba* Miq. Leaves on human cancer cell lines," *Pharmacogn. J.* 5,127–129, (2013)
- T. Jeyalalitha, K. Murugan, M. Umayavalli, "Preliminary phytochemical screening of leaf extracts of *Anthocephalus Cadamba*," *Int. J. Recent Sci. Res.*6, 6608–6611, (2015).
- T. Nishi, K. Yoshizato, et al., "High-efficiency in vivo gene transfer using intra arterial plasmid DNA injection following in vivo electroporation," *Cancer Res.* 56, 1050-1055 (1996).
- U. Zimmermann, & Pilwat, et al., "Dielectric breakdown of cell membranes". *Biophys. J.* 14, 881-899. (1974)
- V. Jothi, V. Raman, R. Sharma, I.Camarillo, and R. Sundararajan, "Electroporation-based Enhanced Anti-Cancer Effect of Veliparib on Triple Negative Breast Cancer Cells," *Proc. Electrostatics Joint Conference.* (2016).
- V. Mlakar, V. Todorovic, M. Cemazar, D. Glavac, G. Sersa, "Electric pulses used in electrochemotherapy and electrogene therapy do not significantly change the expression profile of genes involved in the development of cancer in malignant melanoma cells," *BMC Cancer*, (2009)
- W. Korohoda, M. Grys, Z. Madeja, "Reversible And Irreversible Electroporation Of Cell Suspensions Flowing Through A Localized Dc Electric Field," *Cellular & Molecular Biology Letters*, (2012).
- W. Krassowska, P. D. Filev, "Modeling Electroporation in a Single Cell," *Biophysical Journal*, 92(2), 404–417. (2007).

- W. P. Steward and K. Brown, "Cancer chemoprevention: a rapidly evolving field," *British Journal of Cancer*, vol. 109, no. 1, pp. 1–7, (2013).
- W., Ahmad, F., Ho, W., & Pang, S., "Gc-MS Analysis Of Phytochemical Constituents In Leaf Extracts Of Neolamarckia Cadamba (Rubiaceae) From Malaysia," *International Journal of Pharmacy and Pharmaceutical Sciences*, 6(9), 123-127, (2014).
- Z. Shankayi, S. M. P. Firoozabadi, H. Z. Saraf, "The Endothelial Permeability Increased by Low Voltage and High Frequency Electroporation," *Journal of Biomedical Physics & Engineering*, 3(3), 87–92, (2013).
- Z. Zhou, H. P. He, N. C. Kong, T. J. Wang, X. J. Hao, "Indole alkaloids from the leaves of *Anthocephalus chinensis*," *Helv. Chim. Acta* 91, 2148–2152, (2008).



PTTA UTHM
PERPUSTAKAAN TUNKU TUN AMINAH